



Strain Improvement of *Pseudomonas aeruginosa* F1 for Extracellular L-asparaginase Production Isolated from *Taptapani* a Local Hot Spring in Odisha, India

Biswaprakash Pradhan ¹, Sashi K. Dash ², Chandi C. Rath ³ and Sabuj Sahoo ^{*1}

¹ Pharmaceutical Biotechnology Division, University Department of Pharmaceutical Sciences, Utkal University, Vani Vihar, Bhubaneswar, Odisha-751 004, India

² Department of Seed Technology, Orissa University of Agriculture and Technology, Bhubaneswar, Odisha-751 003, India

³ Deputy Director (Technical), Biotechnology, Govt. of Odisha, Bhubaneswar-751001, India

Received 12 September 2013; accepted in revised form 18 November 2013

Abstract: The purpose of the present investigation is to enhance extracellular L-asparaginase production by subjecting isolated *Pseudomonas aeruginosa* strain F1 (Gene Bank Accession Number: JN412064.1) to improvement by random mutagenesis by ultra-violet (UV) irradiation, Nitrous acid and N-Methyl-N'-nitro-N-nitroso guanidine (NTG) treatment. Mutants were screened as L-asparaginase producers on the basis of enzyme activity in submerged fermentation. UVS-7 mutant obtained after 15-18 minutes of UV exposure showed 1.26 times more enzyme activity (76.6 IU/ml). UVS-7 mutant further mutated by Nitrous acid and UVS7-N-4 mutant obtained after 210-240 Sec Nitrous acid exposure showed 1.58 times increase in enzyme activity (98.41 IU/ml). Then UVS7-N-4 mutant was treated with NTG. UVS7-N-4-N-8 mutant obtained after 210-240 minute NTG exposure showed 2.05 times increase in enzyme activity (124.28 IU/ml) in compared to the isolated wild *Pseudomonas aeruginosa* strain F1(60.64 IU/ml). The results indicated that UV radiation, Nitrous acid and NTG were effective mutagenic agents for strain improvement. Thus these findings have more impact on enzyme economy by enhancing the production of extracellular L-asparaginase for anticancer applications.

Key words: *Pseudomonas aeruginosa* strain F1, mutation, L-asparaginase, strain improvement & Leukemia.

Introduction

L-asparaginase (L-asparagine amidohydrolase, E.C.3.5.1.1) is identified as a potential chemotherapeutic agent since 1961 ¹ and used in combination with other agents in the treatment of acute lymphoblastic leukemia (mainly in children), Hodgkin disease, acute myelocytic leukemia, acute myelomonocytic leukemia, chronic lymphocytic leukemia, lymphosarcoma chemotherapy, pancreatic carcinoma, melanosarcoma, reticulosarcoma ^{2,3} L-asparaginase inhibited the growth of the two human cell lines including

hepatocellular and colon carcinoma ⁴. Preventase (produced from *Aspergillus niger* by DSM Company) and Acrylaway (production from *A. oryzae* by Novozyme Company) are commercially available L-asparaginase that has been used to reduce acrylamide formation in French fries ⁵. L-asparaginase is a model enzyme for the development of new drug delivery system ⁶ and L-asparagine biosensor for leukemia ⁷.

L-asparaginase is an amidohydrolase that catalyzes the hydrolysis of the amino acid L-asparagine to aspartic acid and ammonia. The

*Corresponding author (Sabuj Sahoo)

E-mail: < sabujbiotech@rediffmail.com >

tumor cells have a compromised ability to generate L-asparagine endogenously, either due to low expression levels of asparagine synthetase or insufficient amount of its substrates, aspartate or glutamine⁸. Because of their dependence on exogenous L-asparagine, the cancerous Acute Lymphoblastic Leukemia cells, but not normal cells, can be starved and eliminated by L-asparaginase treatment which depletes the levels of L-asparaginase in circulating pools.

Although almost all living cells produce L-asparaginase, microbial L-asparaginase has received the greater attention because of its apparent advantages in production at large scale in addition to its antitumor property⁹. L-asparaginase has attracted much attention for bulk production with low cost, due to its wide range of applications.

The major motivation for industrial strain development is economic, since the metabolic concentrations produced by wild strains are usually too low for economical process¹⁰. Wild strains frequently produce a mixture of chemically closed related substances. Mutants who synthesize one component as the main product are preferable, since they make possible a simplified process for product recovery¹¹. Changes in the genotype of microorganisms can lead to the biosynthesis of new metabolites with increased yield¹². Genetic variations can be produced by mutation and genetic recombination. The success of strain development depends on an optimal use of mutagenesis (production of mutations) procedures in combination with an effective system for selecting high yielding strain. There were scanty reports on mutational studies for enhancement of L-asparaginase production. Therefore, the present investigation demonstrates the effectiveness of ultra-violet (UV) irradiation¹³, Nitrous acid and N-Methyl-N'-nitro-N-nitrosoguanidine (NTG)¹⁴ in strain improvement for enhanced production of L-asparaginase by *Pseudomonas aeruginosa strain F1*. The mutants were compared with wild type for L-asparaginase production.

Materials and methods

Microorganisms

The thermophilic bacteria *Pseudomonas*

aeruginosa strain F1 isolated from Taptapani hot spring of Odisha, India was employed in the present study. The strain was grown on nutrient agar medium slants at 37°C for two days and used for the study. The slants were sub cultured at monthly intervals and stored at 4°C in the refrigerator.

Cultivation medium and cultural conditions

The composition and of initial cultivation medium was (g/l): L-asparagine- 5.48, yeast extract- 2, beef extract- 2, lactose- 4, Na₂HPO₄·2H₂O - 6.0, KH₂PO₄ - 3.0, NaCl - 0.5, MgSO₄·7H₂O - 0.5, CaCl₂·2H₂O - 0.015 and initial pH was maintained at 6.5^{15,16} with incubation temperature 40°C, inoculum age of 16 h, inoculum size of 6 % w/v, an agitation speed of 150 rpm and 80 % air space. The inoculum was prepared by adding loopful of selected isolates into 50 ml sterilized medium specified above in 250 ml Erlenmeyer (EM) flasks. The flasks were agitated at 150 rpm and incubated at 37°C for 12 h (O.D.₆₀₀ nm=0.6-0.8) in an orbital shaker incubator^{15,16}. After incubation the cells were removed by centrifugation at 6000 × g for 10 min at 4°C. Then the supernatant collected was subjected to assay of extracellular L-asparaginase production.

Enzymatic assay

L-asparaginase activity was measured by direct Nesslerization of ammonia. The activity of L-asparaginase was measured employing the modified method of Wriston^{15,16}.

Preparation of cell suspension

The organism grown on nutrient agar slants were scraped off into sterile water containing tween 80 (1:4000) to give a uniform suspension. The suspension was transferred into a sterile conical flask and thoroughly shaken for 30 min on a rotary shaker. This cell suspension was used for mutation studies.

Mutation and selection

Mutation by UV irradiation

Four mL quantities of cell suspension were pipetted aseptically into sterile petridishes of 100 mm diameter, kept over a flat surface. Then

exposure to UV light was carried out in a "Dispensing Cabinet" fitted with UV lamp TUP 40W germicidal lamp which has about 90 % of its radiation at 2540-2550 Å. The exposure was carried out at a distance of 26.5 cm away from the center of the Germicidal lamp. The exposure times were 0, 3, 6, 9, 12, 15 and 18 min respectively. During the exposure, the lid of the Petridish was removed. Hands were covered with gloves and the plates were gently rotated so as to get uniform exposure to the UV rays. During the exposure all the other sources of light were cut off and the exposure was carried out in dark under red light. The treated cell suspension was transferred into sterile test tubes covered with a black paper and kept in the refrigerator overnight to avoid photo-reactivation.

Each irradiated cell suspension was serially diluted with sterile phosphate buffer. The cell suspension after dilution were plated onto nutrient agar medium and incubated for 24 h at 37°C. The number of colonies in each plate was counted. The percents of number of survivals from each exposure time are shown in Fig. 1. Plates having

less than 1 % survival rate (15-18 min. exposure) were selected for the isolation of mutants. About 12 isolates (UVS1-UVS12) were selected on the basis of macroscopic differences transferred onto nutrient agar slants and tested for their production capacities as described earlier. The best L-asparaginase producing UV mutant strain was selected for further nitrous acid treatment.

Nitrous acid treatment

The cell suspension of the selected strain UVS7 was prepared by using Acetate buffer pH-7. To each 9 ml of the cell suspension in buffer 1 ml sterile stock solution of 0.01 M Sodium Nitrite was added. Sample of 4 ml were withdrawn at 30, 60, 90, 120, 150, 180, 210 and 240 second respectively. Each 1 ml sample was neutralized with 0.5 ml of 0.1M NaOH, serially diluted and plated on the nutrient agar medium.

The colonies from each sample were counted and a percent survival curve was plotted and shown in Fig. 2. Plates having survival rate between 15 and 1 % were selected for the isolation of mutants. The stable mutants UVS-7-




Fig. 1. Effect of UV irradiation on *Pseudomonas aeruginosa* strain F1




Fig. 2. Effect of Nitrous acid on UVS7 *Pseudomonas aeruginosa* strain F1

N-1 to UVS-7-N-10 were selected based on the consistent expression of the phenotypic character up to six generations and maintained on nutrient agar slant. The plates were incubated at 37°C for 24 hours.

NTG treatment and selection of mutants

The Nitrous acid mutants of UVS-7-N-4 maintained on nutrient agar slants was used for NTG treatment. The selected Nitrous acid treated mutant (UVS-7-N-4) was sub-cultured onto nutrient agar and incubated at 37°C for 24 h. The cell suspension was prepared as described earlier. The cells were suspended in sterile phosphate buffer and shaken to get a uniform suspension. The NTG (100 mg) was accurately weighed and dissolved in 25 mL of phosphate buffer at a temperature of 40°C to minimize decomposition and sterilized by passing through sterile bacterial filter (0.22 µm). Ten mL of the cell suspension was added to 25 mL of NTG solution and immediately incubated at 37°C in a water bath. At appropriate time intervals 5 mL samples were withdrawn and centrifuged immediately. The supernatant was decanted, and then the cell pellet was washed with sterile distilled water and finally resuspended in 5 mL of sterile distilled water. A total of seven samples were withdrawn from the cell suspension each in 5 mL quantities after exposure to NTG for 30, 60, 90, 120, 150 and 180 min respectively. A control also was included without exposure to NTG.

The above treated samples were serially diluted and 100 µL of each sample was spread onto NA plates by spread plate technique. The plates were

incubated at 37°C for 24 h. The colonies from each plate were counted and a percent survival curve was plotted and shown in Fig. 3. Plates having less than 1 % survival rate were selected for the isolation of mutants. Colonies were selected on the basis of macroscopic characteristics. A total of 12 colonies were picked up from the plates and transferred onto nutrient agar slants. These were labeled as *UVS7-N-4-N-1* to *UVS7-N-4-N-12*. They were incubated at 37°C for 24 h to get good growth. All the selected mutants were tested separately for their L-asparaginase production capacity.

Statistical analysis

All values presented here are the average values of triplicate analysis.

Results and discussion

The wild strain of *Pseudomonas aeruginosa strain F1* was subjected to strain improvement programme with a view to obtain increased L-asparaginase production and to achieve greater stability of the organism. The first method chosen was UV irradiation.

Isolation of UV Mutants and their L-asparaginase Activity

The selected mutants were provided with the code number *UVS1* to *UVS12* and these mutants were tested for their L-asparaginase producing capabilities and Results are prescribed in Table 1. The results indicated that among UV-mutants, *UVS7* was found to be the highest L-asparaginase producer (76.6 IU/mL) and it was 1.26 times




Fig. 3. Effect of NTG on UVS7-N-4 *Pseudomonas aeruginosa strain F1*

Table 1. L-asparaginase production capacity of UV mutants

UV mutants	L-asparaginase activity (IU/mL)
UVS1	67.16
UVS2	52.84
UVS3	72.32
UVS4	68.41
UVS5	55.31
UVS6	59.87
UVS7	76.60
UVS8	49.18
UVS9	68.23
UVS10	74.17
UVS11	54.33
UVS12	68.28
<i>Pseudomonas aeruginosa strain F1</i>	60.40

higher than the parent strain (60.4 IU/mL). The mutant, *UVS7* was selected for further strain improvement studies using Nitrous acid.

Selection of Nitrous acid Mutants and their L-asparaginase Activity

The selected mutant *UVS7* was subjected to nitrous acid treatment. Mutation frequency was observed to be high when the survival rates were between 15 and 1 %. The dose survival curve was plotted for selecting the mutants between 15 % and 1 % survivals. Plates having survival rate between 15 and 1 % (210 and 240 sec) were selected for the isolation of mutants. The survival curve was plotted by taking exposure time

(Second) on X-axis and % survival on Y-axis (Fig 2.). A total of 10 mutants were selected and determined for their L-asparaginase production capacities by submersed fermentation, the activity was determined and presented in Table 2. Out of 10 mutants *UVS7-N-4* showed maximum L-asparaginase activity of 98.41 IU/mL, which was 1.58 times more than the wild isolate *Pseudomonas aeruginosa strain F1*(62.2 IU/mL).

Selection of NTG Mutants and their L-asparaginase Activity

The selected UV & Nitrous acid-mutant, *UVS7-N-4* was subjected to NTG treatment. A total of 12 mutants were selected and they were provided

Table 2. L-asparaginase production capacity of Nitrous acid mutants

Nitrous acid Mutants	L-asparaginase activity (IU/mL)
UVS7-N-1	57.16
UVS7-N-2	92.84
UVS7-N-3	79.32
UVS7-N-4	98.41
UVS7-N-5	59.31
UVS-N-6	79.27
UVS-N-7	76.43
UVS-N-8	63.18
UVS-N-9	88.23
UVS-N-10	54.17
<i>Pseudomonas aeruginosa strain F1</i>	62.20

with the code numbers *UVS7-N-4-N-1* to *UVS7-N-4-N-12*. These were evaluated for their L-asparaginase production capacities. The results are shown in Table 3. The results indicated that the mutant *UVS7-N-4-N-8* was the highest L-asparaginase yielding strain (124.28 IU/mL). Thus the NTG treatment resulted a mutant that produced 2.05 times higher L-asparaginase than the parent wild strain (60.64 IU/mL), which is a significant increase in yield. From the results, it is evident that UV, Nitrous acid and NTG were effective mutagenic agents for strain improvement of *Pseudomonas aeruginosa strain F1*. By employing these techniques, a superior mutant (*UVS7-N-4-N-8*) with a productivity of 2.05 times higher yield than the parent strain was obtained.

Summary

L-asparaginase is an enzyme used as an antitumor agent in the treatment of various types of leukemia. Earlier, this enzyme was produced by several bacteria but for commercial production only mutants of *E. coli* were used. The shortage of this enzyme, subsequently lead to the discovery of novel organisms producing L-asparaginase.

Here we have used our thermophilic isolate *Pseudomonas aeruginosa strain F1*, obtained from Taptapani hot spring of Odisha, India for the enhanced production of L-asparaginase. There are many evidences on the successful employment of strain improvement techniques to increase enzyme yield in various industrially important microorganisms^{17,18,19}.

A stable mutant of *Pseudomonas aeruginosa strain F1* for L-asparaginase production was obtained through an efficient and safe mutation method by using Physical (UV radiation) and chemical (Nitrous acid & N-methyl-N-nitro-N-nitrosoguanidine) mutagens. The effectiveness of the mutagens helped us to achieve a 2.05 times enhanced production of L-asparaginase in the mutated strain than the wild strain that helps in minimizing the cost of extracellular L-asparaginase production for its biotechnological applications.

Acknowledgements

The funding support by the All India Council for Technical Education, New Delhi, India through "Research Promotion Scheme" is gratefully acknowledged.

Table 3. L-asparaginase production capacity of NTG mutants

NTG Mutants	L-asparaginase activity (IU/mL)
<i>UVS7-N-4-N-1</i>	86.76
<i>UVS7-N-4-N-2</i>	113.84
<i>UVS7-N-4-N-3</i>	92.62
<i>UVS7-N-4-N-4</i>	88.41
<i>UVS7-N-4-N-5</i>	105.31
<i>UVS7-N-4-N-6</i>	99.27
<i>UVS7-N-4-N-7</i>	76.63
<i>UVS7-N-4-N-8</i>	124.28
<i>UVS7-N-4-N-9</i>	78.36
<i>UVS7-N-4-N-10</i>	107.76
<i>UVS7-N-4-N-11</i>	84.43
<i>UVS7-N-4-N-12</i>	68.86
<i>Pseudomonas aeruginosa strain F1</i>	60.64

References

1. **Broome, J.D. (1961).** Evidence that the L-asparaginase activity of guinea pig serum is responsible for its antilymphoma effects. *Nature.*, 191: 1114-1115.
2. **Chakrabarti, R. (1997).** L-asparaginase perspectives on the mechanisms of action and resistance.

- Int. J. Pediatr. Hematol. Oncol. 4: 597-611.
3. **Gulati, R., Saxena, R.K., Gupta, R. (1997).** A rapid plate assay for screening L-asparaginase producing micro-organisms. *Lett. Appl. Microbiol.* 24: 23-26.
 4. **Oza, V.P., Parmar, P.P., Kumar, S., Subramanian, R.B. (2010).** Anticancer properties of highly purified L-asparaginase from *Withaniasomnifera* L. against acute lymphoblastic leukemia. *Appl. Biochem. and Biotech* 160(6): 1833-40.
 5. **Lapmak, K., Lumyong, S., Thongkuntha, S., Wongputtisin, P., Sardsud, U. (2010).** L-asparaginase production by *Bipolaris* sp. BR438 isolated from brown rice in Thailand. *Chiang Mai. J. Sci.* 37(1): 160-164.
 6. **Teodor, E., Litescu, S.C., Lazar, V., Somoghi, R. (2009).** Hydrogel-magnetic nanoparticles with immobilized L-asparaginase for biomedical applications. *J. Mater. Sci. Mater. Med.* 20: 1307-1314.
 7. **Verma, N., Kumar, K., Kaur, G., Anand, S. (2007).** *E. coli* K-12 asparaginase-based asparagine biosensor for leukemia. *Artif Cells Blood Substit. Immobil. Biotechnol.* 35: 449-456.
 8. **Aslanian, A.M., Kilberg, M.S. (2001).** Multiple adaptive mechanisms affect asparagine synthetase substrate availability in asparaginase-resistant MOLT-4 human leukaemia cells. *Biochem. J.* 358: 59-67.
 9. **Kavitha, A., Vijayalakshmi, M. (2012).** A Study on L-Asparaginase of *Nocardia levis* MK-VL 113. *The Scientific World Journal*, 1-5.
 10. **Lin, C., Yang, L., Xu, G., Wu, J. (2011).** Enhancement of Haloacetate Dehalogenase Production by Strain Mutation and Condition Optimization. *Biotechnology and Bioprocess Engineering* 16: 923-929.
 11. **Khushoo, A., Pal, Y., Singh, B.N., Mukherjee, K.J. (2004).** Extracellular expression and single step purification of recombinant *Escherichia coli* L-asparaginase II. *Protein Expression and Purification* 38: 29-36.
 12. **Zambare, V. (2012).** Strain improvement of alkaline protease from *Trichoderma reesei* MTCC-3929 by physical and chemical mutagen. *Research: Bio-Engineering* 1(1): 25-28.
 13. **Hungund, B.S., Gupta, S.G. (2010).** Strain improvement of *Gluconacetobacter xylinus* NCIM 2526 for bacterial cellulose production. *African Journal of Biotechnology* 9(32): 5170-5172.
 14. **Krishna, E.R., Kumar, P.S., Kumar, B.V. (2011).** Strain improvement of selected strain *Bacillus subtilis* (MTCC No.10619) for enhanced production of antimicrobial metabolites. *J. Microbiol. Biotech. Res.* 1(3): 32-38.
 15. **Pradhan, B., Dash, S.K., Sahoo, S. (2013).** Optimization of some physical and nutritional parameters for the production of L-asparaginase by isolated thermophilic *Pseudomonas aeruginosa* strain F1. *Biosciences Biotechnology Research Asia* 10(1): 1-7.
 16. **Pradhan, B., Dash, S.K., Sahoo, S. (2013).** Optimization of some physical and nutritional parameters for the production of L-asparaginase by isolated thermophilic *Bacillus subtilis* strain *hswx88*. *Pharma nest* 4(4): 627-636.
 17. **Lee, M.Y., Hong, K.N., Soon, D.K., Witoon, P. (2007).** Quality of chungkukjangs prepared with various *Bacillus* strains. *Int. J. Food Sci. Tech.* 42(5): 587-592.
 18. **Mocktar, C., Govinden, U., Sturm, A.W., Essack, S.Y. (2007).** TEM-145 and TEM-146 β -lactamases produced by *Escherichia coli* isolates from state hospitals in KwaZulu-Natal, South Africa. *Afr. J. Biotech.* 6: 493-496.
 19. **Wang, H.Y., Liu, D.M., Liu, Y., Cheng, C.F., Ma, Q.Y., Huang, Q., Zhang, Y.Z. (2007).** Screening and mutagenesis of a novel *Bacillus pumilus* strain producing alkaline protease for dehairing. *Lett. Appl. Microbiol.* 44(1): 1-6.